## **PIROXANTRONE**

#### NSC - 349174

## **Chemical Name:**

Anthra[1,9-cd]pyrazol-6(2H)-one, 5-[3-(aminopropyl)amino]-7,10-dihydroxy-2-[2-[(2-hydroxyethyl)amino]ethyl]-, dihydrochloride

#### Other Names:

Anthrapyrazole derivative; Oxantrazole

CAS Registry Number: 105118-12-5

Molecular Formula: C<sub>21</sub>H<sub>25</sub>N<sub>5</sub>O<sub>4</sub>·2HCl M.W.: 484.4

## **Approximate Solubility:** (mg/mL)

H <sub>2</sub> O	> 11.20
Buffer, pH4	> 9.60
Buffer, pH9	> 11.50
Ethanol	< 0.73

DMA	0.70
DMSO	> 9.60
CHCl <sub>3</sub>	< 0.77
EtOAc	< 0.71
t-BuOH	< 0.71

## Stability:

#### **Bulk:**

Samples stored at 25 °C under both light and dark conditions at 50 °C in the dark for three months showed no significant degradation (HPLC).

#### **Solution:**

A solution containing 10.2 mg/mL in water showed approximately 5% degradation in 24 hours. The test sample exhibited < 10% degradation in 48 hours.

## **Ultraviolet Absorption:**

 $(H_2O)$ 

λ

∧ <sub>max</sub>	ε
254 ± 2nm	$10,358 \pm 500$
$295 \pm 2nm$	$6,146 \pm 400$
$318 \pm 2$ nm	$6,889 \pm 400$
$333 \pm 2$ nm	$6,677 \pm 400$
$406 \pm 2$ nm	$4,827 \pm 400$
$515 \pm 2$ nm	$16,592 \pm 600$

# High Performance Liquid Chromatography:

Column:

Zorbax C<sub>8</sub>, 250 x 4.6 mm i.d.,

30 x 4.6 mm i.d.

**Mobile Phase:** 

5% CH<sub>3</sub>CN/20% DMF /75% (0.2M

ammonium acetate/L mobile phase,

adjusted with acetic acid to pH 4.5.)

Flow Rate:

1 mL/min

**Detection:** 

UV at 254 nm

Sample Preparation:

10 mg of sample diluted to 10 mL with

internal standard solution.

**Internal Standard:** 

1.0 mg/mL benzamide in water

**Retention Volume:** 

11.14 mL (NSC-349174)

6.17 mL (I.S.)